

سوال (b) لا

[3] Duality

$$\text{if } g(t) \Longleftrightarrow G(f)$$

$$\text{then } G(t) \Longleftrightarrow g(-f)$$

Ex Find F.T of $AT \text{sinc}(tT)$

using Duality




$$A \text{rect}\left(\frac{t}{T}\right) \Longleftrightarrow AT \text{sinc}(fT)$$

$$AT \text{rect}\left(\frac{t}{T}\right) \Longleftrightarrow A \text{rect}\left(\frac{-f}{T}\right)$$



$$-f=0, f=0$$

* في حالة ال rect فيكون متساوي (-)

كلاهما واحد



Limited ~~Free~~ in time \Longleftrightarrow unlimited Free



* using Duality

$$1) A \text{rect}\left(\frac{t}{\tau}\right) \Longleftrightarrow A\tau \text{sinc}(f\tau)$$

$$2) A\tau \text{sinc}(\tau t) \Longleftrightarrow A \text{rect}\left(\frac{f}{\tau}\right)$$

$$A \sin(2\omega t) \Longleftrightarrow \frac{A}{2\omega} \text{rect}\left(\frac{f}{2\omega}\right)$$

$$\tau = 2\omega$$

[EX] Find F.T of $\sin(mt)$

using Duality

$$A \text{rect}(t/\tau) \Longleftrightarrow A\tau \text{sinc}(f\tau)$$

$$A\tau \text{sinc}(t\tau) \Longleftrightarrow A \text{rect}(f/\tau)$$

$$\text{sinc}(mt) \Longleftrightarrow \frac{1}{m} \text{rect}\left(\frac{f}{m}\right)$$

$$\tau = m, A\tau = 1, A = \frac{1}{m}$$

[2] sec 4

[4] Time shift Property

$$\text{If } x(t) \xrightarrow{\quad\quad\quad} G(f)$$

$$\text{then } x(t \pm t_0) \xrightarrow{\quad\quad\quad} G(f) \cdot e^{\pm j 2 \pi f t_0}$$

نفس الـ t_0


Ex: Find F.T for $x(t) = A \text{rect}(t - t_0)$

amp \swarrow

Sol

using time shift

$$A \text{rect}\left(\frac{t}{\tau}\right) \xrightarrow{\quad\quad\quad} A\tau \underbrace{\sin(f\tau)}_{G(f)}$$




$$A \text{rect}\left(\frac{t - t_0}{\tau}\right) \xrightarrow{\quad\quad\quad} A\tau \sin(f\tau) \cdot e^{-j 2 \pi f t_0}$$

Ex Find F.T For $g(t) = A \text{rect}\left(\frac{t - \frac{\tau}{2}}{\tau}\right)$

Sol

using time shift

$$A \text{rect}\left(\frac{t - \frac{\tau}{2}}{\tau}\right) \Rightarrow$$



$$A \text{rect}\left(\frac{t - \frac{\tau}{2}}{\tau}\right) \Rightarrow A \tau \text{sinc}(\tau f) \cdot e^{-j2\pi f \frac{\tau}{2}}$$


$$A \tau \text{sinc}(\tau f) \cdot e^{-j\pi f \tau}$$

Ex Find F.T of $g(t)$ as shown.

$$g(t) = A \text{rect}\left(\frac{t - \tau/2}{\tau}\right)$$

$$- A \text{rect}\left(\frac{t + \tau/2}{\tau}\right)$$

$$- A \text{rect}$$



using Linearity & time shift

$$A \text{rect}\left(\frac{t - \tau/2}{\tau}\right) \Longleftrightarrow A\tau \text{sinc}(f\tau) \cdot e^{-j2\pi f \frac{\tau}{2}}$$

$$A \text{rect}\left(\frac{t + \tau/2}{\tau}\right) \Longleftrightarrow A\tau \text{sinc}(f\tau) \cdot e^{j2\pi f \frac{\tau}{2}}$$

$$G(f) = A\tau \text{sinc}(f\tau) \left[e^{-j\pi f\tau} - e^{+j\pi f\tau} \right]$$

$$\therefore G(f) = (-2j) A\tau \text{sinc}(f\tau) \cdot \sin(\pi f\tau) \quad \cdot * \frac{-2j}{-2j}$$

5 Frequency-shift property

$$\text{if } g(t) \Longleftrightarrow G(f)$$

$$\text{then } g(t) \cdot e^{\pm j2\pi f_0 t} \Longleftrightarrow G(f \mp f_0)$$

5

sec 4

Ex Find F.T of $A \text{rect}\left(\frac{t}{\tau}\right) \cdot e^{-j2\pi f_c t}$

Sol

using Freq. shift

$$A \text{rect}\left(\frac{t}{\tau}\right) \xLeftrightarrow AT \text{sinc}(f\tau)$$

$$G(f) = AT \text{sinc}(f\tau)$$

$$G(f + f_c) = AT \text{sinc}(\tau(f + f_c))$$

Ex
7.6 Find F.T of

$$x(t) = A \text{rect}\left(\frac{t}{\tau}\right) \cdot \cos(2\pi f_c t)$$

$$x(t) = A \text{rect}\left(\frac{t}{\tau}\right) \left[\frac{e^{j2\pi f_c t} + e^{-j2\pi f_c t}}{2} \right]$$



$$= \frac{1}{2} \left[A \text{rect}\left(\frac{t}{\tau}\right) \cdot e^{j2\pi f_c t} + A \text{rect}\left(\frac{t}{\tau}\right) \cdot e^{-j2\pi f_c t} \right]$$

using superposition & Linearity


$$A \text{rect}\left(\frac{t}{\tau}\right) \cdot e^{j2\pi f_c t} \xLeftrightarrow AT \text{sinc}((f + f_c)\tau)$$

$$A \operatorname{rect}\left(\frac{t}{\tau}\right) \cdot e^{-j2\pi f_c t} \xLeftrightarrow A\tau \cdot \operatorname{sinc}((f+f_c)\tau)$$

$$\therefore G(f) = \frac{1}{2} \left[A\tau \cdot \operatorname{sinc}((f-f_c)\tau) + A\tau \operatorname{sinc}((f+f_c)\tau) \right]$$




$$g(t) = A \operatorname{rect}\left(\frac{t}{\tau}\right) \cdot \cos(2\pi f_c t)$$




$$\cos(2\pi f_c t)$$

$g(t)$



$$G(f) \propto \frac{1}{2} \left[A\tau \operatorname{sinc}((f-f_c)\tau) + A\tau \operatorname{sinc}((f+f_c)\tau) \right]$$



* Modulation Theory:-

$$m(t) \cdot \cos(2\pi f_c t) \iff \frac{1}{2} \left[M(f+f_c) + M(f-f_c) \right]$$

$$m(t) \cdot \frac{e^{j\theta} - e^{-j\theta}}{2j} \iff \frac{1}{2j} \left[M(f-f_c) - M(f+f_c) \right]$$

[6] Area under the curve $g(t)$

$$\text{Area} = \int_{-\infty}^{\infty} g(t) \cdot dt$$

$$G(f) \propto \int_{-\infty}^{\infty} g(t) \cdot e^{-j2\pi f t} \cdot dt$$

[8] Sec 4


$$\therefore \text{Area} = G(0)$$

Ex Find Area of $g(t) = A \text{ rect}(\frac{t}{\tau})$

Sol

$$\text{Area} = A * \tau$$

$$\text{Area} = A\tau$$



another solution

$$G(f) = A\tau \text{ sinc}(f\tau)$$

$$\text{Area} = G(0) = A\tau \text{ sinc}(0) = A\tau$$

[7] Area under Curve $G(f)$

$$\text{Area} = \int_{-\infty}^{\infty} G(f) \cdot df$$

I.F.t

$$g(t) = \int_{-\infty}^{\infty} G(f) \cdot e^{j2\pi f \cdot t} \cdot df$$

[9] Sec 4


$$\Rightarrow t=0$$

$$\therefore \text{Area} \propto \omega(0)$$

EX Find Area under Curve

$$G(f) \propto \frac{1}{\sqrt{2\pi f}}$$

$$|G(f)| \propto \frac{1}{\sqrt{1+4\pi^2 f^2}}$$



Q.1

Report 2

Find F.T For

[1] $q(t) \propto 3 \sin(t-3)$

[2] $q(t) \propto \text{rect}\left(\frac{t+1.5}{7}\right)$

[3] $q(t) \propto 3 \cdot e^{-|t-2|}$

[4] $\sec 4$

$$q(t) \propto e^{-t} \cdot u(t)$$

$$\therefore \text{Area} \propto \omega(0)$$

$$\text{Area} \propto 1 \cdot \frac{1}{2} = \frac{1}{2}$$